

UNIVERSITY OF OKLAHOMA
GRADUATE COLLEGE

NON-LOCAL THERMODYNAMIC EQUILIBRIUM
SPECTRUM SYNTHESIS OF
TYPE Ia SUPERNOVAE

A DISSERTATION
SUBMITTED TO THE GRADUATE FACULTY
in partial fulfillment of the requirements for the
degree of
DOCTOR OF PHILOSOPHY

By
PETER E. NUGENT
Norman, Oklahoma
1997

**NON-LOCAL THERMODYNAMIC EQUILIBRIUM
SPECTRUM SYNTHESIS OF
TYPE Ia SUPERNOVAE**

**A DISSERTATION APPROVED FOR THE
DEPARTMENT OF PHYSICS AND ASTRONOMY**

BY

Edward Baron (Chair)

David Branch

John Cowan

William Romanishin

Susan Postawko

©Copyright by Peter Nugent 1997

All Rights Reserved.

Acknowledgements

I have many people to thank for getting me to where I am today. Without their help this dissertation would have never been realized. First on the list are my parents, who showed me the importance of an education (though I don't think they had 24 years in mind when they sent me off to kindergarten) and gave me the freedom to pursue whatever I wanted to in school. I also want to thank my brother James for being there for me.

I wish to thank my grandfather for giving me my first lessons in electricity and magnetism. My curiosity in physics started in his basement workshop sifting through resistors and capacitors and playing with the oscilloscope. He also gave me my first telescope, something which started off as a hobby and led to my chosen profession of astronomy.

I thank Professors Thomas Bross and E.O. LaCasce. They saw me through high school and college physics respectively. The skills they taught me and the work ethic they instilled in me has served me well the past seven years of graduate school and will continue to do so for many years to come.

I thank Ed Baron and David Branch for introducing me to supernovae and advising me throughout graduate school. Their teaching, patience, generosity and friendship has been greatly appreciated. I look forward to many future collaborations.

Finally I wish to thank the two most important people in my life. Anya, whose cute little smile when I come home from a hard day at work makes it all worthwhile, and Deepa, whose unfailing love and encouragement these past seven years has meant everything to me. I wouldn't have made it this far without you.

Contents

1	Introduction	1
1.1	NLTE Spectrum Synthesis and SNe Ia	1
1.2	Phoenix	3
2	SNe Ia 1992A and 1981B	4
2.1	Abstract	4
2.2	Introduction	4
2.3	Description of the Models	5
2.4	Comparison to Other Work	6
2.5	Model Parameters	8
2.6	Results	9
2.7	Discussion	12
2.8	Conclusions	15
3	Physics and the Hubble Constant	17
3.1	Abstract	17
3.2	Introduction	17
3.3	Theory	19
3.4	Hydrodynamical Models	20
3.5	Spectrum Synthesis	21
3.6	Results	22
3.7	Conclusions	24
4	A Spectroscopic Sequence Among SNe Ia	26
4.1	Abstract	26
4.2	Introduction	26
4.3	Observations	28
4.4	Theory	33
4.5	Discussion and Conclusions	37
5	Hydrodynamic Models of SNe Ia	41
5.1	Abstract	41
5.2	Introduction	42

5.3	Calculations	43
5.4	Results	46
5.4.1	Model W7	46
5.4.2	Sub-Chandrasekhar Models	50
5.5	Conclusions	61
6	NLTE vs. LTE	65
6.1	Abstract	65
6.2	Differences	65
6.3	Line Identifications	68
A	γ-ray Transport in Type Ia Supernovae	83
B	α — The Thermalization Parameter	89

List of Tables

2.1	The parameters used to fit SN 1992A and 1981B with the resulting absolute magnitudes and colors.	12
4.1	Observed $\mathcal{R}(\text{Si II})$ and $\mathcal{R}(\text{Ca II})$	30
4.2	Synthetic $\mathcal{R}(\text{Si II})$ and $\mathcal{R}(\text{Ca II})$	36
5.1	Magnitudes and Colors of W7 Models. M_{Bol} , M_B , and M_V are the bolometric, B and V absolute magnitudes of the models respectively. $B - V$ and $U - B$ are the associated colors.	46
5.2	Magnitudes and Colors of the Sub-Chandrasekhar Models. M_{Bol} , M_B , and M_V are the bolometric, B and V absolute magnitudes of the models respectively. $B - V$ and $U - B$ are the associated colors. . . .	54
5.3	4 Most Abundant Species for Each Model at 20 Days After Explosion. The relative concentrations by number of the four most abundant species (in parts per million) for a particular zone (labeled by its velocity). The models W7, WW2, and LA4 have $\eta = 1.0, 1.1, 1.4$, respectively.	61
6.1	Magnitudes for the NLTE and LTE models.	66
B.1	The color and magnitude differences for a set of models with different values of α	92

List of Figures

2.1	Observed and synthetic spectra of SN 1992A.	10
2.2	Observed and synthetic spectra of SN 1981B.	11
2.3	Two synthetic spectra with differing density profiles.	13
3.1	The luminosity of Type Ia supernovae as a function of rise-time. . . .	23
4.1	A plot of the synthetic (left) and observed (right) spectral sequences.	29
4.2	Observed M_B vs. $\mathcal{R}(\text{Si II})$ and $\mathcal{R}(\text{Ca II})$	31
4.3	Photometry in U , B , V , R and I for the synthetic spectra.	33
4.4	The colors for the synthetic spectra.	38
4.5	$\sqrt{\kappa_{RM}}$ for all the models at two different values of τ	39
5.1	The synthetic spectrum for the W7 model at 20 d past explosion . . .	47
5.2	The W7 model at 23 d ($\eta = 1.3$) compared with an observed spectrum of SN 1992A	49
5.3	The time evolution of the W7 model near maximum light	51
5.4	Model WW2 15 d after explosion for three choices of η	52
5.5	Model LA4 15 d after explosion for three choices of η	53
5.6	The Sobolev optical depths of the helium lines	55
5.7	The Sobolev optical depths throughout the atmosphere of Model WW2	57
5.8	The synthetic spectra of WW2 ($\eta = 1.1$) and LA4 ($\eta = 1.5$) at 20 d compared with the observed spectrum of SN 1994D	58
5.9	The synthetic spectra of WW2 ($\eta = 1.1$) and LA4 ($\eta = 1.5$) at 20 d compared with the observed spectrum of SN 1991bg	59
5.10	The density and temperature profiles for WW2 and W7	63
5.11	The Sobolev optical depth of the Co II $^1D - ^3G^\circ$ $\lambda 2605.2$ line	64
6.1	A comparison of LTE and NLTE spectra.	67
6.2	The percentage difference in temperature between the NLTE and LTE spectra.	69
6.3	The b_i 's for the NLTE model.	70
6.4	The Sobolev optical depths for O I and C I throughout the line forming region in the atmosphere.	72

6.5	The Sobolev optical depths for Na I and Mg II throughout the line forming region in the atmosphere.	73
6.6	The Sobolev optical depths for Si II and S II throughout the line forming region in the atmosphere.	74
6.7	The Sobolev optical depths for Ca II and Ti II throughout the line forming region in the atmosphere.	75
6.8	The Sobolev optical depths for Fe II and Co II throughout the line forming region in the atmosphere.	76
A.1	Geometric lay-out for the γ -ray deposition calculation.	85
A.2	Geometric lay-out for the analytic γ -ray deposition function.	86
A.3	A Graph of the semi-analytic deposition function vs. the numerical absorption calculation.	87
A.4	The 20 day deposition functions of models W7, WW2 and LA4. . . .	88
B.1	ϵ as a function of wavelength for W7 at 20 days after explosion. . . .	94
B.2	A good fit to the spectrum of SN 1992A for $\alpha = 0.1$	95
B.3	Unacceptable fits to the spectrum of SN 1992A for $\alpha = 0.01$ and 0.3 . . .	96
B.4	The synthetic spectrum for the W7 model at 23 d past explosion . . .	97